

Ryanodine Receptor 2 (RYR2) (dephoSer2808) pAb serum

Quality Control Certificate of Analysis

Catalogue No.: A010-35AP

Unit Size: 100 µl Lot No.: 0611-05

Background: The ryanodine receptor (RyR2) is a Ca²⁻ channel of cardiac muscle that plays a central role in EC coupling. The binding of Ca²⁻ to RyR2 opens the channel and Ca²⁻ stored in the SR moves through the channel into the cytosol to initiate muscle contraction (Bers, 2002). Abnormal structure and function of ryanodine receptors has been reported in failing hearts, with Ser-2809 phosphorylation appearing elevated in clinical situations which may contribute to the abnormal Ca²⁻ handling characteristics of cardiac muscle in these conditions (Wehrens and Marks, 2003). Serine-2809 can be phosphorylated by PKA or CaMKII (Rodriguez *et al.*, 2003), which is coincident with significant change in RyR2 channel function typified by an increased open probability (Witcher *et al.*, 1991; Valdivia *et al.*, 1995; Marx *et al.*, 2000), the abrogation of the inhibitory effects of CaM (Witcher *et al.*, 1991) and Mg²⁻ (Hain *et al.*, 1995), dissociation of regulatory factors, expression of subconductance states and the expression of channel activity at diastolic Ca²⁻ concentrations (Marx *et al.*, 2000). This antibody recognises Dephosphorylated Ser-2808 (human sequence) or Ser-2809 from rabbit.

Description: Affinity purified **Rabbit** polyclonal antibody (A010-35) to dephosphorylated ryanodine receptor Ser-2809.

Immunogen: Synthetic peptide (Y₂₈₀₂NRTRRISQT₂₈₁₁) corresponding to amino acids surrounding the dephosphorylated serine residue at position 2808 of RyR2, which was conjugated to keyhole limpet hemocyanin (KLH) by carbodiimide cross linkage.

Antibody Isotype: IgG.

Antibody Purity: Protein A Affinity Purified.

Specificity: The antibody recognises dephosphorylated Serine-2808 of the ryanodine receptor. Binding of the antibody to its target epitope is blocked in the presence of a phosphopeptide containing the dephospho Ser-2808 epitope.


Species Cross Reactivity: Reacts with dephos Ser-2808 of phospholamban from canine, rat and sheep species.


Vial Constituents: Lyophilised affinity purified A010-35AP Ab (100 µl) in 0.1M Tris-citrate pH 7.4 with 20%v/v stabiliser solution.

Storage Instructions: Lyophilised antibody is stable at 4°C when stored with desiccant. Reconstitute lyophilised powder in 100 μ I of 18 M Ω H $_2$ O, aliquot and store frozen at -80°C for 1 year. Avoid freeze - thaw cycles.

Tested Applications: WB 1:500. Not yet tested in other applications, therefore, optimal dilutions/concentrations should be determined by the user.

PO₃H₂Specific





A010-35AP : lot 0611-05 (1:500) 2 μ l in 1 ml of Tris buffer + 5% milk powder. 16 hours at 4°C

Secondary antibody = Goat anti Rabbiut HRP Chemilluminescent detection: 60 second exposure

WB using 1:500 A010-35AP against PKA, CaMKII treated and untreated canine cardiac Sarcoplasmic Reticulum. 7 % gel PVDF membrane. Specific band at 350 kDa. Additional band at 80 - 85 kDa.

Related Products: RyR2 dephospho Ser-2809 epitope peptide (P010-35); RYR2 Phospho Ser-2809 antibody (A010-30AP).

Background References:

- Bers, D. M. (2002): Nature 415, 198-205.
- Hain, J., Onoue, H., Mayrleitner, M., Fleischer, S., and Schindler, H. (1995): J Biol Chem 270, 2074-81.
- Marx, S. O., Reiken, S., Hisamatsu, Y., Jayaraman, T., Burkhoff, D., Rosemblit, N., and Marks, A. R. (2000): Cell 101, 365-76.
- Rodriguez, P., Bhogal, M. S., and Colyer, J. (2003): J Biol Chem 278, 38593-600.
- Valdivia, H. H., Kaplan, J. H., Ellis-Davies, G. C., and Lederer, W. J. (1995): Science 267, 1997-2000.
- Wehrens, X. H., and Marks, A. R. (2003): *Trends Biochem Sci* **28**, 671-8.
- Witcher, D. R., Kovacs, R. J., Schulman, H., Cefali, D. C., and Jones, L. R. (1991): J Biol Chem 266, 11144-52.